

Before you start, assemble the supplies you will need for the injection:


- medication
- 1 cc (ml) syringe
- alcohol swab
- gauze

IMPORTANT: Use a new syringe and needle every day

To administer the medication you will have to prepare the medication, select and prepare the injection site, then finally inject the medication. **Begin by washing your hands.**

A. To Prepare the Medication:


1. Remove the protective caps from the two vial of medication and use the alcohol swab to clean the rubber covering the vial. Allow to dry.



2. Take the syringe out of its sterile wrapper, and remove the protective cap.

3. Pull back on the syringe plunger (inner part) and draw up ____cc (ml) units of air.

4. Insert the needle into the vial of medication and inject the air into the bottle.




5. Keep the needle in the bottle and turn the bottle upside down. Hold the bottle straight up and confirm that the needle tip is submerged in liquid. Withdraw **at least ____cc(ml) = ____units** of medication- take more than you need because you will always have some air bubbles mixed in that you will need to squirt back into the bottle.

6. With the needle still in the bottle, hold the bottle firmly and flick the syringe hard so that the bubbles move up to the surface. Then push on the plunger so that the bubbles and the extra fluid move up into the bottle. Keep pushing on the plunger until you have exactly **cc(ml) = ____units of medication** remaining in the syringe. Withdraw the needle from the bottle and recap the needle to protect it until you are ready to give the injection.

B.To Select the Injection Site: (Select new site each day)

You will be injecting the medication deep into the subcutaneous tissue, which is the fat just beneath the skin. There are three major sites where you can inject:

- the back of the upper arm
- the abdomen (except for navel and waist)
- the outer side of the thigh (not too close to the groin or knee)



Be sure to avoid veins, moles, and scars.

C. To Inject the Medication:

1. Swab the skin with the alcohol swab and allow the area to dry.

2. Uncap the needle and grasp the syringe like a pencil. With your other hand, pinch a fold of skin.


3. Gently insert the needle at a 90° angle straight into the skin. Be sure to insert the needle fully.



4. Release your pinch and slowly inject the medication by pushing the plunger all the way.

5. Wait 5 seconds and withdraw the needle, keeping the same angle as during the insertion.

6. Apply gauze and firm pressure to the injection site until any bleeding has stopped.



D. Discard the Used Syringe and Needles:

Discard the syringe into a plastic container that has a lid (you do not have to recap the needle before you discard it). Return the covered container to a pharmacy or the clinic.

THIS INFORMATION IS INTENDED AS A GUIDELINE AND MAY NOT CONTAIN INFORMATION ABOUT ALL ASPECTS OF CARE